

Bibliography (cont.)

- Coffin PO, Rowe C, Oman N, et al. Illicit opioid use following changes in opioids prescribed for chronic non-cancer pain among persons living with and without HIV. *AIDS 2020: 23rd International AIDS Conference Virtual*. July 6-10, 2020. Abstract PEB0347.
- Cofrancesco J Jr, Scherzer R, Tien PC, et al. Illicit drug use and HIV treatment outcomes in a US cohort. *AIDS*. 2008;22:237-245.
- Collaboration of Observational HIV Epidemiological Research Europe (COHERE) Study Group, Sabin CA, Smith CJ, d'Arminio Monforte A, et al. Response to combination antiretroviral therapy: Variation by age. *AIDS*. 2008;22:1463-1473.
- Connor EM, Sperling RS, Gelber R, et al. Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. *N Engl J Med*. 1994;331:1173-1180.
- Courlet P, Livio F, Guidi M, et al. Polypharmacy, drug-drug interactions, and inappropriate drugs: New challenges in the aging population with HIV. *Open Forum Infect Dis*. 2019;6(12):ofz531.
- Cuzin L, Katlama C, Cotte L, et al. Ageing with HIV: Do comorbidities and polymedication drive treatment optimization? *HIV Med*. 2017;18(6):395-401.
- DeFrancesco D, Wit FW, Bürkle, et al. Do people living with HIV experience greater age advancement than their HIV-negative counterparts? *AIDS*. 2019;33(2):259-268.
- DeLorenze GN, Weisner C, Tsai AL, et al. Excess mortality among HIV-infected patients diagnosed with substance use dependence or abuse receiving care in a fully integrated medical care program. *Alcohol Clin Exp Res*. 2011;35:203-210.
- Dickover RE, Dillon M, Gillette SG, et al. Rapid increases in load of human immunodeficiency virus correlate with early disease progression and loss of CD4+ cells in vertically infected infants. *J Infect Dis*. 1994;170:1279-1284.
- Dorenbaum A, Cunningham CK, Gelber RD, et al. Two-dose intrapartum/newborn nevirapine and standard antiretroviral therapy to reduce perinatal HIV transmission: A randomized trial. *JAMA*. 2002;288:189-198.