

参考文献

- Aboud M, et al. Dolutegravir Versus Ritonavir-Boosted Lopinavir Both With Dual Nucleoside Reverse Transcriptase Inhibitor Therapy in Adults With HIV-1 Infection in Whom First-Line Therapy Has Failed (DAWNING): An Open-Label, Non-Inferiority, Phase 3b Trial. *Lancet Infect Dis*. 2019 Mar;19(3):253-264.
- Aboud M, Orkin C, Podzamczar D, et al. Durable suppression 2 years after switch to DTG + RPV 2-drug regimen: SWORD 1&2 studies. 22nd IAS Conference: July 23-27, 2018. Amsterdam, The Netherlands. Abstract THPEB047.
- Aboud M, Orkin C, Podzamczar D, et al. Efficacy and safety of dolutegravir-rilpivirine for maintenance of virological suppression in adults with HIV-1: 100-week data from the randomised, open-label, phase 3 SWORD-1 and SWORD-2 studies. *Lancet HIV*. 2019;6(9):e576-e587.
- Arribas JR, DeJesus E, van Lunzen J, et al. Simplification to single-tablet regimen of elvitegravir, cobicistat, emtricitabine, tenofovir DF from multi-tablet ritonavir-boosted protease inhibitor plus coformulated emtricitabine and tenofovir DF regimens: week 96 results of STRATEGY-PI. *HIV Clin Trials*. 2017;18(3):118-125.
- Arribas JR, Girard PM, Landman R, et al. Dual treatment with lopinavir-ritonavir plus lamivudine versus triple treatment with lopinavir-ritonavir plus lamivudine or emtricitabine and a second nucleos(t)ide reverse transcriptase inhibitor for maintenance of HIV-1 viral suppression (OLE): a randomised, open-label, non-inferiority trial. *Lancet Infect Dis*. 2015;15(7):785-792.
- Arribas JR, Pialoux G, Gathe J, et al. Simplification to coformulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus continuation of ritonavir-boosted protease inhibitor with emtricitabine and tenofovir in adults with virologically suppressed HIV (STRATEGY-PI): 48 week results of a randomised, open-label, phase 3b, non-inferiority trial. *Lancet Infect Dis*. 2014;14(7):581-589.
- Baxter JD, Mayers DL, Wentworth DN, et al. A randomized study of antiretroviral management based on plasma genotypic antiretroviral resistance testing in patients failing therapy. *AIDS*. 2000;14:F83-F93.
- Borghetti A, Baldin G, Lombardi F, et al. Efficacy and tolerability of lamivudine plus dolutegravir as a switch strategy in a multicentre cohort of patients with suppressed HIV-1 replication. *HIV Med*. 2018;19(7):452-454.